

**SECTION-I**

**2. Write short answers to any EIGHT (8) questions: (16)**

**(i) Define implicit function.**

**Ans** If  $x$  and  $y$  are so mixed up and  $y$  cannot be expressed in terms of the independent variable  $x$ , then  $y$  is called an implicit function. For example,

$$1. x^2 + xy + y^2 = 0$$

$$2. \frac{xy^2 - y + 9}{xy} = + 1 \text{ are implicit functions of } x \text{ and } y.$$

Symbolically, it is written as  $f(x, y) = 0$ .

**(ii)  $f(x) = 3x^4 - 2x^2$ ,  $g(x) = \frac{2}{\sqrt{x}}$ , find  $f(g(x))$ .**

$$\text{Ans} \quad f(x) = 3x^4 - 2x^2 \quad g(x) = \frac{2}{\sqrt{x}}$$

$$\begin{aligned} fog(x) &= f[g(x)] = 3\left(\frac{2}{\sqrt{x}}\right)^4 - 2\left(\frac{4}{\sqrt{x}}\right)^2 \\ &= 3\left(\frac{16}{x^2}\right) - 2\left(\frac{16}{x}\right) = \frac{48}{x^2} - \frac{32}{x} \\ &= \frac{48 - 32x}{x^2} = \frac{8(6 - x)}{x^2} \end{aligned}$$

**(iii) Evaluate  $\lim_{x \rightarrow 2} \frac{\sqrt{x} - \sqrt{2}}{x - 2}$ .**

$$\begin{aligned} \text{Ans} &= \lim_{x \rightarrow 2} \frac{\sqrt{x} - \sqrt{2}}{x - 2} \times \frac{\sqrt{x} + \sqrt{2}}{\sqrt{x} + \sqrt{2}} \\ &= \lim_{x \rightarrow 2} \frac{x - 2}{(x - 2)(\sqrt{x} + \sqrt{2})} \\ &= \frac{1}{\sqrt{2} + \sqrt{2}} = \frac{1}{2\sqrt{2}} \end{aligned}$$

**(iv) Find derivative by definition of  $x^2$ .**

$$\text{Ans} \quad f(x) = x^2$$

1.  $f(x + \delta x) = (x + \delta x)^2$
2.  $f(x + \delta x) - f(x) = (x + \delta x)^2 - x^2$   
 $= x^2 + 2x \delta x + (\delta x)^2 - x^2$   
 $= 2x \delta x + (\delta x)^2 = (2x + \delta x) \delta x$
3.  $\frac{f(x + \delta x) - f(x)}{\delta x} = \frac{(2x + \delta x) \delta x}{\delta x} = 2x + \delta x \quad (\delta x \neq 0)$
4.  $\lim_{\delta x \rightarrow 0} \frac{f(x + \delta x) - f(x)}{\delta x}$   
 $\lim_{\delta x \rightarrow 0} (2x + \delta x) = 2x$   
*i.e.,*  $f'(x) = 2x$

(v) Differentiate w.r.t. 'x'  $\sqrt{\frac{a-x}{a+x}}$ .

**Ans** Let  $y = \sqrt{\frac{a-x}{a+x}}$  and  $u = \frac{a-x}{a+x}$

Then  $y = u^{1/2}$

$$\text{Now, } \frac{dy}{du} = \frac{1}{2} u^{-1/2} = \frac{1}{2} u^{-1/2}$$

$$\text{and } \frac{du}{dx} = \frac{d}{dx} \left[ \frac{a-x}{a+x} \right]$$

$$\begin{aligned} &= \frac{\left[ \frac{d}{dx}(a-x) \right] (a+x) - (a-x) \left[ \frac{d}{dx}(a+x) \right]}{(a+x)^2} \\ &= \frac{(0-1)(a+x) - (a-x)(0+1)}{(a+x)^2} = \frac{-a-x-a+x}{(a+x)^2} \\ &= \frac{-2a}{(a+x)^2} \end{aligned}$$

Using the formula,

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}, \quad \text{we have}$$

$$\begin{aligned} \frac{d}{dx} \left( \sqrt{\frac{a-x}{a+x}} \right) &= \frac{1}{2} u^{-1/2} \left[ \frac{-2a}{(a+x)^2} \right] \\ &= \frac{1}{2} \left( \frac{a-x}{a+x} \right)^{1/2} \times \frac{-2a}{(a+x)^2} \quad \left( \because u = \frac{a-x}{a+x} \right) \end{aligned}$$

$$= \frac{(a-x)^{-1/2}}{(a+x)^{-1/2}} \times \frac{-a}{(a+x)^2} = \frac{-a}{(a-x)^{1/2} (a+x)^{3/2}}$$

(vi) Find  $\frac{dy}{dx}$ , if  $x^2 - 4xy - 5y = 0$ .

**Ans**  $x^2 - 4xy - 5y = 0$

Differentiating the equation w.r.t x:

$$\frac{d}{dx}(x^2 - 4xy - 5y) = 0$$

$$\frac{d}{dx}x^2 - \frac{d}{dx}(4xy) - \frac{d}{dx}(5y) = 0$$

$$2x - 4\left[x \frac{d}{dx}(y) + y \frac{d}{dx}(x)\right] - 5 \frac{d}{dx}(y) = 0$$

$$2x - 4\left(x \frac{dy}{dx} + y\right) - 5 \frac{dy}{dx} = 0$$

$$2x - 4x \frac{dy}{dx} - 4y - 5 \frac{dy}{dx} = 0$$

$$-\frac{dy}{dx}(4x + 5) = 4y - 2x$$

$$\frac{dy}{dx} = -\frac{(4y - 2x)}{4x + 5}$$

$$= \frac{2x - 4y}{4x + 5} = \frac{2(x - 2y)}{4x + 5}$$

(vii) Prove that  $\frac{d}{dx}(\cot^{-1} x) = -\frac{1}{1+x^2}$ .

**Ans** Let  $y = \cot^{-1} x \Rightarrow y = \frac{1}{\cot x}$

$$\therefore \cot y = x \quad (i)$$

Differentiating w.r.t. x,

$$\frac{d}{dx}(\cot y) = \frac{d}{dx}(x)$$

$$-\operatorname{cosec}^2 y \cdot \frac{dy}{dx} = 1$$

$$\frac{dy}{dx} = -\frac{1}{\operatorname{cosec}^2 y}$$

$$= \frac{1}{1 + \cot^2 y} = -\frac{1}{1 + x^2} \quad \text{Proved.}$$

(viii) Find  $\frac{dy}{dx}$ , if  $y = x \cos y$ .

**Ans**

$$y = x \cos y$$

Differentiate w.r.t.  $x$ .

$$\frac{dy}{dx} = \frac{d}{dx}(x \cos y)$$

$$= x \frac{d}{dx}(\cos y) + \cos y \frac{d}{dx}(x)$$

$$= x \left( -\sin y \cdot \frac{dy}{dx} \right) + \cos y (1)$$

$$= -x \sin y \frac{dy}{dx} + \cos y$$

$$= \frac{dy}{dx} + x \sin y \frac{dy}{dx} = \cos y = \frac{dy}{dx} (1 + x \sin y) = \cos y$$

$$\frac{dy}{dx} = \frac{\cos y}{1 + x \sin y}$$

(ix) Find  $f'(x)$ , if  $f(x) = \sqrt{\ln(e^{2x} + e^{-2x})}$ .

**Ans**

$$\text{Let } u = e^{2x} + e^{-2x}$$

Then  $f(x) = (\ln u)^{1/2}$  and

$$\begin{aligned} f'(x) &= \frac{d}{dx} (\ln u)^{1/2} = \frac{d}{du} (\ln u)^{1/2} \times \frac{du}{dx} = \left[ \frac{1}{2} (\ln u)^{1/2-1} \frac{d}{du} (\ln u) \right] \times \frac{d}{dx} (e^{2x} + e^{-2x}) \\ &= \left( \frac{1}{2} \cdot \frac{1}{(\ln u)^{1/2}} \cdot \frac{1}{u} \right) \cdot (e^{2x} \cdot 2 + e^{-2x} \cdot (-2)) \\ &= \frac{1}{2} \cdot \frac{1}{\sqrt{\ln(e^{2x} + e^{-2x})}} \cdot \frac{1}{e^{2x} + e^{-2x}} \cdot 2(e^{2x} - e^{-2x}) \\ &= \frac{e^{2x} + e^{-2x}}{\sqrt{\ln(e^{2x} + e^{-2x})} \times (e^{2x} + e^{-2x})} \end{aligned}$$

(x) Find  $y_2$ , if  $x = at^2$ ,  $y = bt^4$ .

**Ans**

$$x = at^2$$

;

$$y = bt^4$$

$$\frac{dx}{dt} = \frac{d}{dt}(at^2)$$

;

$$\frac{dy}{dt} = \frac{d}{dt}(bt^4)$$

$$= 2at$$

;

$$= 4bt^3$$

By Chain Rule:

$$\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx}$$

$$= 4bt^3 \times \frac{1}{2at}$$

$$\therefore y_1 = \frac{2bt^2}{a}$$

$$\frac{d^2y}{dx^2} = \frac{2b}{a} \cdot \frac{d}{dx}(t^2)$$

$$\begin{aligned}\therefore y_2 &= \frac{2b}{a} \cdot (2t) \cdot \frac{d}{dx}(t) \\ &= \frac{4bt}{a} \cdot \frac{dt}{dx} \\ &= \frac{4bt}{a} \times \frac{1}{2at} = \frac{2b}{a^2}\end{aligned}$$

(xi) Define Maclaurin series.

**Ans** We have  $a_0 = f(0)$ ,  $a_1 = f'(0)$ ,  $a_2 = f''(0) \frac{(0)}{2!}$

$$a_3 = \frac{f'''(0)}{3!}, a_4 = \frac{f''''(0)}{4!}$$

Following the above pattern, we can write

$$a_n = \frac{f^{(n)}(0)}{n!}$$

Thus substituting these values in the power series, we have

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots$$

This expansion of  $f(x)$  is called the Maclaurin series.

(xii) Determine the interval in which  $f(x)$  is increasing or decreasing if  $f(x) = \sin x$ ,  $x \in (0, \pi)$ .

**Ans**  $f(x) = \sin x$   
 $f'(x) = \cos x$

$$f'(x) \text{ is } +\text{ve } \forall x \in (0, \frac{\pi}{2})$$

$\therefore f(x)$  is increasing.

$$f'(x) = \cos x \text{ is } -\text{ve for all } x \in (\frac{\pi}{2}, \pi)$$

then  $f(x)$  is decreasing.

### 3. Write short answers to any EIGHT (8) questions: (16)

(i) Using differential, find  $\frac{dy}{dx}$  when  $xy - \ln x = c$ .

**Ans**  $xy - \ln x = c$

Taking differential on both sides,

$$d(xy - \ln x) = d(c)$$

$$x dy + y dx - \frac{1}{x} dx = 0$$

$$x dy + y dx - \frac{1}{x} dx = 0$$

$$x dy = \frac{dx}{x} - y dx$$

$$= dx \left( \frac{1}{x} - y \right)$$

$$\frac{dy}{dx} = \frac{\frac{1}{x} - y}{x} = \frac{1 - xy}{x^2}$$

(ii) Evaluate  $\int \frac{(\sin x + \cos^2 x)}{\cos^2 x \cdot \sin x} dx$ .

**Ans**

$$\begin{aligned} &= \int \left( \frac{\sin x}{\cos^2 x \cdot \sin x} + \frac{\cos^2 x}{\cos^2 x \cdot \sin x} \right) dx \\ &= \int \left( \frac{1}{\cos^2 x} + \frac{1}{\sin x} \right) dx \\ &= \int \sec^2 x dx + \int \operatorname{cosec} x dx \\ &= \tan x + \ln |\operatorname{cosec} x - \cot x| + c \end{aligned}$$

(iii) Find  $\int x(\sqrt{x+1}) dx ; x > 0$ .

**Ans**

$$\begin{aligned} &\int x \cdot x^{1/2} dx + \int x dx \\ &\int x^{3/2} dx + \frac{x^2}{2} + c \\ &= \frac{x^{5/2}}{\frac{5}{2}} + \frac{x^2}{2} + c \\ &= \frac{2x^{5/2}}{5} + \frac{x^2}{2} + c \end{aligned}$$

(iv) Evaluate  $\int a^{x^2} \cdot x dx ; a > 1$ .

**Ans** Put  $x^2 = t$ , then  $x dx = \frac{1}{2} dt$

$$\text{Thus } \int a^{x^2} \cdot x dx = \int a^t \times \frac{1}{2} dt$$

$$= \frac{1}{2} \int a^t dt = \frac{1}{2} \frac{a^t}{\ln a} + c = \frac{a^{x^2}}{2 \ln a} + c$$

(v) Find the anti-derivative of  $x \cdot e^x$ .

**Ans**  $\int x \cdot e^x dx$  Let  $u = x$  and  $dv = e^x dx$

Then  $du = 1 \cdot dx$  and  $v = e^x$

Applying the formula for integration by parts, we have

$$\begin{aligned}\int x e^x dx &= x e^x - \int e^x \times 1 dx \\ &= x e^x - e^x + c\end{aligned}$$

(vi) Evaluate  $\int e^x (\cos x + \sin x) dx$ .

**Ans**  $= \int e^x \cos x dx + \int e^x \sin x dx$

Integrating 1<sup>st</sup> integral by parts,

$$\begin{aligned}&= e^x \int \cos x dx - \int \left[ \frac{d}{dx}(e^x) \cdot \int \cos x dx \right] dx + \int e^x \sin x dx + c \\ &= e^x \cdot \sin x - \int e^x \cdot \sin x dx + \int e^x \sin x dx + c \\ &= e^x \sin x + c\end{aligned}$$

(vii) State 'Fundamental Theorem' of calculus.

**Ans** If  $f$  is continuous on  $[a, b]$  and  $\phi'(x) = f(x)$ , that is,  $\phi(x)$  is any anti-derivative of  $f$  on  $[a, b]$ , then

$$\int_a^b f(x) dx = \phi(b) - \phi(a)$$

Note that the difference  $\phi(b) - \phi(a)$  is independent of the choice of anti-derivative of the function  $f$ .

(viii) Compute  $\int_{-1}^1 (x^{1/3} + 1) dx$ .

$$\begin{aligned}\text{Ans} \quad \int_{-1}^1 (x^{1/3} + 1) dx &= \int_{-1}^1 x^{1/3} dx + \int_{-1}^1 1 dx \\ &= \left[ \frac{x^{4/3}}{\frac{4}{3}} \right]_{-1}^1 + [x]_{-1}^1 \\ &= \frac{3}{4} [(1)^{4/3} - (-1)^{4/3}] + [(1) - (-1)] \\ &= \frac{3}{4} [1 - 1] + 2 = 2\end{aligned}$$

(ix) Find the area above x-axis and under the curve  $y = 5 - x^2$  from  $x = -1$  to  $x = 2$ .

**Ans**

$$\begin{aligned} \text{Area} &= \int_{-1}^2 (5 - x^2) dx \\ &= 5 \int_{-1}^2 dx - \int_{-1}^2 x^2 dx \\ &= 5[x]_{-1}^2 - \left[ \frac{x^3}{3} \right]_{-1}^2 \\ &= 5[2 - (-1)] - \frac{1}{3}[8 - (-1)^3] \\ &= 5[3] - \frac{1}{3}[8 + 1] \\ &= 15 - \frac{1}{3}(9) = 15 - 3 = 12 \text{ sq. units} \end{aligned}$$

(x) Solve the differential equation  $\sin y \cdot \operatorname{cosec} x \cdot \frac{dy}{dx} = 1$ .

**Ans**

$$\sin y \operatorname{cosec} x \frac{dy}{dx} = 1$$

$$\sin y dy = \frac{dx}{\operatorname{cosec} x}$$

$$\sin y dy = \sin x dx$$

$$\int \sin y dy = \int \sin x dx + c'$$

$$-\cos y = -\cos x + c'$$

$$\cos y = \cos x - c'$$

$$\cos y = \cos x + c$$

(xi) Define 'decision variables'.

**Ans** The variables used in the system of linear inequalities relating to the problems of everyday life are non-negative constraints. These non-negative constraints play an important role for taking decision. So these variables are decision variables.

(xii) Graph solution set of inequality  $2x + y \geq 2$  in  $x - y$  plane.

**Ans** We draw the graph of the line  $2x + y = 2$  joining the points  $(1, 0)$  and  $(0, 2)$ . The point  $(0, 0)$  does not satisfy the inequality  $2x + y > 0$  because  $2(0) + 0 = 0 \not> 2$ . Thus the graph of the inequality  $2x + y \geq 2$  is the closed half not on the origin-side of the line  $2x + y = 2$ .

The intersection point  $(2, -2)$  can be found by solving the equation  $2x + y = 2$ .

**4. Write short answers to any NINE (9) questions: (18)**

- (i) Find the coordinates of the point that divides the join of  $A(-6, 3)$  and  $B(5, -2)$  internally in ratio  $2 : 3$ .

**Ans** Here  $k_1 = 2$ ,  $k_2 = 3$ ,  $x_1 = -6$ ,  $x_2 = 5$

By the formula, we have  $y_1 = 3$ ,  $y_2 = -2$

$$x = \frac{(2 \times 5) + 3(-6)}{2 + 3} = \frac{10 - 18}{5} = \frac{-8}{5}$$

$$\text{and } y = \frac{(2(-2)) + 3 \times 3}{2 + 3} = \frac{-4 + 9}{5} = \frac{5}{5} = 1$$

Coordinates of the required points are  $\left(\frac{-8}{5}, 1\right)$ .

- (ii) Find the slope and inclination of the line joining the points  $A(-2, 4)$  and  $B(5, 11)$ .

**Ans** Let  $A(-2, 4)$  and  $B(5, 11)$

$$\text{Slope of } AB = m = \frac{11 - 4}{5 - (-2)} = \frac{7}{5 + 2} = \frac{7}{7} = 1$$

$$m = \tan \theta = 1 \Rightarrow \text{inclination} = \theta = \tan^{-1}(1) = 45^\circ$$

- (iii) By means of slopes show that points  $A(-1, -3)$ ,  $B(1, 5)$  and  $C(2, 9)$  are collinear.

**Ans** We know that the points  $A$ ,  $B$  and  $C$  are collinear, if the line  $AB$  and  $BC$  have the same slopes.

$$\text{Slope of } AB = m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{5 - (-3)}{1 - (-1)} = \frac{5 + 3}{1 + 2} = \frac{8}{2} = 4$$

$$\text{and Slope of } BC = \frac{9 - 5}{2 - 1} = \frac{4}{1} = 4$$

$$\therefore \text{Slope of } AB = \text{Slope of } BC$$

Thus  $A$ ,  $B$  and  $C$  are collinear.

- (iv) Find equation of the line through  $(-4, 7)$  and parallel to the line  $2x - 7y + 4 = 0$ .

**Ans**  $2x - 7y + 4 = 0$

$$-7y = -2x - 4$$

$$7y = 2x + 4$$

$$\frac{7y}{7} = \frac{2x}{7} + \frac{4}{7}$$

$$y = \frac{2}{7}x + \frac{4}{7}$$

∴ Slope of required linear  $= m = \frac{2}{7}$

Equation of line through  $(-4, 7)$  with  $m = \frac{2}{7}$  is

$$y - y_1 = m(x - x_1)$$

$$y - 7 = \frac{2}{7}(x - (-4))$$

$$y - 7 = \frac{2}{7}(x + 4)$$

$$7y - 49 = 2x + 8$$

$$2x - 7y = 49 + 8 \Rightarrow 2x - 7y = 57$$
$$\Rightarrow 2x - 7y - 57 = 0$$

(v) Find equation of circle with centre at  $(5, -2)$  and radius 4.

Ans Here  $c(h, k) = c(5, -2)$  and  $r = 4$

Equation of required circle is

$$(x - h)^2 + (y - k)^2 = r^2$$

$$(x - 5)^2 + (y - (-2))^2 = (4)^2$$

$$(x - 5)^2 + (y + 2)^2 = (4)^2$$

$$x^2 + 25 - 10x + y^2 + 4y + 4 = 16$$

$$x^2 + y^2 - 10x + 4y + 29 = 16$$

$$x^2 + y^2 - 10x + 4y + 29 - 16 = 0$$

$$x^2 + y^2 - 10x + 4y + 13 = 0$$

(vi) Find focus and vertex of the parabola  $y^2 = -8(x - 3)$ .

Ans  $y^2 = -8(x - 3) \Rightarrow y^2 = -8x$  where  $x = x - 3$

Comparing it with  $y^2 = -4ax$

$$-4a = -8$$

$$4a = 8$$

$$a = \frac{8}{4} = 2$$

Coordinates of focus:

$(-a, 0)$

$$x = -a, y = 0$$

$$x - 3 = -2$$

$$x = 3 - 2 = 1$$

$f(1, 0)$

Coordinates of vertex,  $(0, 0)$

$$x = 0, y = 0$$

$$x - 3 = 0$$

$v(0, 0)$

(vii) Find equation of tangent to the parabola  $x^2 = 16y$  at the point whose abscissa is 8.

**Ans** Since  $x = 8$  lies on the parabola,

Substituting this value of  $x$  into the given equation, we find

$$x^2 = 16y$$

$$(8)^2 = 16y$$

$$64 = 16y \Rightarrow y = 4$$

Thus we have to find equations of tangent and normal at  $(8, 4)$ .

Slope of the tangent to the parabola at  $(8, 4)$  is 1. An equation of the tangent to the parabola at  $(8, 4)$  is

$$y - 4 = x - 8$$

$$x - y + 4 - 8 = 0$$

$$x - y - 4 = 0$$

Slope of the normal at  $(8, 4)$  is  $-1$ . Therefore, equation of the normal at the given point is

$$y - 4 = -(x - 8)$$

$$y - 4 = +8 - x$$

$$x + y - 4 - 8 = 0$$

$$x + y - 12 = 0$$

(viii) Find foci and vertices of the ellipse  $25x^2 + 9y^2 = 225$ .

**Ans**

$$25x^2 + 9y^2 = 225$$

Dividing both sides by 225

$$\frac{25x^2}{225} + \frac{9y^2}{225} = \frac{225}{225}$$

$$\frac{x^2}{9} + \frac{y^2}{16} = 1$$

$$\text{Here, } a^2 = 25 \Rightarrow a = \pm 5$$

$$b^2 = 9 \Rightarrow b = \pm 3$$

$$e^2 = \frac{a^2 - b^2}{a^2} = \frac{25 - 9}{25} \Rightarrow \frac{16}{25} \Rightarrow e = \frac{4}{5}$$

$$ae = 5\left(\frac{4}{5}\right) = 4$$

$$\frac{a}{e} = \frac{5}{\frac{4}{5}} = \frac{5 \times 5}{4} = \frac{25}{4}$$

$$f = (0, \pm ae) = (0, \pm 4)$$

$$V(0, \pm a) = V(0 \pm 5)$$

(ix) Find the angle between the vectors  $\underline{u} = 2\mathbf{i} - \mathbf{j} + \mathbf{k}$  and  $\underline{v} = -\mathbf{i} + \mathbf{j}$ .

**Ans**

$$\underline{u} = 2\mathbf{i} - \mathbf{j} + \mathbf{k}$$

$$\underline{v} = -\mathbf{i} + \mathbf{j}$$

$$\underline{u} \cdot \underline{v} = (2\mathbf{i} - \mathbf{j} + \mathbf{k}) \cdot (-\mathbf{i} + \mathbf{j} + 0\mathbf{k})$$

$$= (2)(-1) + (-1)(1) + (1)(0) = -3.$$

$$|\underline{u}| = |2\mathbf{i} - \mathbf{j} + \mathbf{k}| = \sqrt{(2)^2 + (-1)^2 + (1)^2} = \sqrt{6}$$

$$\text{and } |\underline{v}| = |-\mathbf{i} + \mathbf{j} + 0\mathbf{k}| = \sqrt{(-1)^2 + (1)^2 + (0)^2} = \sqrt{2}$$

$$\cos \theta = \frac{\underline{u} \cdot \underline{v}}{|\underline{u}| \cdot |\underline{v}|}$$

$$\cos \theta = \frac{-3}{\sqrt{6} \sqrt{2}} = \frac{-\sqrt{3}}{2}$$

$$\therefore \theta = \frac{5\pi}{6}$$

(x) Find scalar  $\alpha$  so that the vectors  $2\mathbf{i} + \alpha\mathbf{j} + 5\mathbf{k}$  and  $3\mathbf{i} + \mathbf{j} + \alpha\mathbf{k}$  are perpendicular.

**Ans**

$$\text{Let } \underline{u} = 2\mathbf{i} + \alpha\mathbf{j} + 5\mathbf{k}$$

$$\text{and } \underline{v} = 3\mathbf{i} + \mathbf{j} + \alpha\mathbf{k}$$

It is given that  $\underline{u}$  and  $\underline{v}$  are perpendicular.

$$\therefore \underline{u} \cdot \underline{v} = 0$$

$$\Rightarrow (2\mathbf{i} + \alpha\mathbf{j} + 5\mathbf{k}) \cdot (3\mathbf{i} + \mathbf{j} + \alpha\mathbf{k}) = 0$$

$$\Rightarrow 6 + \alpha + 5\alpha = 0$$

$$6\alpha = -6$$

$$\alpha = \frac{-6}{6} = -1$$

$$\therefore \alpha = -1$$

(xi) If  $\underline{v}$  is a vector for which  $\underline{v} \cdot \mathbf{i} = 0$ ,  $\underline{v} \cdot \mathbf{j} = 0$ ,  $\underline{v} \cdot \mathbf{k} = 0$ , find  $\underline{v}$ .

**Ans**

$$\text{Let } \underline{v} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$$

(i)

$$\therefore \underline{v} \cdot \mathbf{i} = 0$$

$$\therefore (a\mathbf{i} + b\mathbf{j} + c\mathbf{k})(\mathbf{i} - 0\mathbf{j} + 0\mathbf{k}) = 0$$

$$(a)(1) + (b)(0) + (c)(0) = 0$$

$$a = 0$$

(ii)

$$\therefore \underline{v} \cdot \mathbf{j} = 0$$

$$\therefore (a\mathbf{i} + b\mathbf{j} + c\mathbf{k}) \cdot (0\mathbf{i} + \mathbf{j} + 0\mathbf{k}) = 0$$

$$\begin{aligned}
 \therefore (\underline{a})(0) + b(1) + c(0) &= 0 \\
 \therefore b &= 0 \quad (\text{iii})
 \end{aligned}$$

$$\begin{aligned}
 \therefore \underline{v} \cdot \underline{k} &= 0 \\
 \therefore (\underline{a}\underline{i} + \underline{b}\underline{j} + \underline{c}\underline{k}) \cdot (\underline{0}\underline{i} + \underline{0}\underline{j} + \underline{k}) &= 0 \\
 \therefore (\underline{a} \times \underline{0}) + (\underline{b} \times \underline{0}) + (\underline{c} \times \underline{1}) &= 0 \\
 \therefore c &= 0 \quad (\text{iv})
 \end{aligned}$$

Putting (ii), (iii) and (iv) in eq. (i),

$$\begin{aligned}
 \underline{v} &= (0)\underline{i} + 0(\underline{j}) + 0(\underline{k}) \\
 \underline{v} &= 0 \quad (\text{Zero / Null vector})
 \end{aligned}$$

(xii) Prove that  $\underline{a} \times (\underline{b} + \underline{c}) + \underline{b} \times (\underline{c} + \underline{a}) + \underline{c} \times (\underline{a} + \underline{b}) = 0$ .

**Ans**

$$\begin{aligned}
 \underline{a} \times \underline{b} + \underline{a} \times \underline{c} + \underline{b} \times \underline{c} + \underline{b} \times \underline{a} + \underline{c} \times \underline{a} + \underline{c} \times \underline{b} &= 0 \\
 \underline{a} \times \underline{b} + \underline{a} \times \underline{c} + \underline{b} \times \underline{c} - \underline{a} \times \underline{b} - \underline{a} \times \underline{c} - \underline{b} \times \underline{c} &= 0 \\
 &= 0 \text{ R.H.S}
 \end{aligned}$$

(xiii) Find the value of  $\alpha$ , so that  $\alpha\underline{i} + \underline{j}$ ,  $\underline{i} + \underline{j} + 3\underline{k}$  and  $2\underline{i} + \underline{j} - 2\underline{k}$  are coplanar.

**Ans** Let  $\underline{u} = \alpha\underline{i} + \underline{j}$ ,  $\underline{v} = \underline{i} + \underline{j} + 3\underline{k}$  and  
 $\underline{w} = 2\underline{i} + \underline{j} - 2\underline{k}$

Triple scalar product

$$\begin{aligned}
 [\underline{u} \underline{v} \underline{w}] &= \begin{bmatrix} \alpha & 1 & 0 \\ 1 & 1 & 3 \\ 2 & 1 & -2 \end{bmatrix} \\
 &= \alpha(-2 - 3) + -1(-2 - 6) + 0(1 - 2) \\
 &= -5\alpha + 8
 \end{aligned}$$

The vectors will be coplaner if

$$-5\alpha + 8 = 0$$

$$-5\alpha = -8$$

$$\alpha = \frac{8}{5}$$

## SECTION-II

**NOTE:** Attempt any Three (3) questions.

Q.5.(a) If  $f(x) = \begin{cases} 3x & \text{if } x \leq -2 \\ x^2 - 1 & \text{if } -2 < x < 2 \\ 3 & \text{if } x \geq 2 \end{cases}$  (5)

discuss continuity at  $x = 2$  and  $x = -2$ .

**Ans** (i) At  $x = 2$

- (a)  $f(2) = 3$  (Defined)
- (b) L.H.L. =  $\lim_{x \rightarrow 2^-} (x^2 - 1) = (2)^2 - 1 = 3$
- R.H.L. =  $\lim_{x \rightarrow 2^+} (3) = 3$
- $\therefore$  L.H.L. = R.H.L.
- (c)  $\lim_{x \rightarrow 2} f(x) = 3$
- $\therefore \lim_{x \rightarrow 2} f(x) = f(2)$

Hence  $f(x)$  is continuous at  $x = 2$ .

- (ii) At  $x = -2$
- (a)  $f(-2) = 3(-2) = -6$  (Defined)
- (b) L.H.L. =  $\lim_{x \rightarrow -2^-} 3x = 3(-2) = -6$
- R.H.L. =  $\lim_{x \rightarrow -2^+} (x^2 - 1) = (-2)^2 - 1 = 3$

$\therefore$  L.H.L.  $\neq$  R.H.L.

Hence  $f(x)$  is discontinuous at  $x = -2$ . There is no need to investigate (c).

- (b) If  $y = e^x \sin x$ , show that  $\frac{d^2y}{dx^2} - 2 \frac{dy}{dx} + 2y = 0$ . (5)

**Ans**

$$y = e^x \sin x$$

(i)

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(e^x \sin x) \\ &= e^x \frac{d}{dx}(\sin x) + \sin x \frac{d}{dx}(e^x) \\ &= e^x \cos x + e^x \sin x \\ &= e^x (\sin x + \cos x)\end{aligned}$$

(ii)

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx}[e^x (\sin x + \cos x)] \\ &= e^x \frac{d}{dx}(\sin x + \cos x) + (\sin x + \cos x) \frac{d}{dx} e^x \\ &= e^x (\cos x - \sin x) + e^x (\sin x + \cos x) \\ &= e^x \cos x - e^x \sin x + e^x \sin x + e^x \cos x \\ &= 2e^x \cos x\end{aligned}$$

(iii)

From (ii),

$$\begin{aligned}\frac{dy}{dx} &= e^x \cos x + e^x \sin x \\ &= e^x \cos x + y\end{aligned} \quad (\text{From (i)})$$

$$e^x \cos x = \frac{dy}{dx} - y$$

Putting (iv) in (iii),

$$\frac{d^2y}{dx^2} = 2 \left( \frac{dy}{dx} - y \right)$$

$$= 2 \frac{dy}{dx} - 2y$$

$$\frac{d^2y}{dx^2} - 2 \frac{dy}{dx} + 2y = 0 \quad \text{Proved.}$$

Q.6.(a) Integrate  $\int \frac{12}{x^3 + 8} dx$ . (5)

**Ans**  $\frac{12}{x^3 + 8} = \frac{12}{(x)^3 + (2)^3} = \frac{12}{(x+2)(x^2 - 2x + 4)}$

Making partial fractions,

$$\frac{12}{(x+2)(x^2 - 2x + 4)} = \frac{A}{x+2} + \frac{Bx+C}{x^2 - 2x + 4} \quad (i)$$

$$12 = A(x^2 - 2x + 4) + (Bx + C)(x + 2) \quad (ii)$$

For A, let  $x + 2 = 0$

$$x = -2$$

Putting  $x = -2$  in eq (ii),

$$12 = A((-2)^2 - 2(-2) + 4) + 0$$

$$12 = A(4 + 4 + 4) + 0$$

$$12 = 12A$$

$$\frac{12}{12} = A$$

$$\boxed{1 = A}$$

Expanding eq. (ii),

$$\begin{aligned} 12 &= Ax^2 - 2Ax + 4A + Bx^2 + 2Bx + Cx + 2C \\ &= (A + B)x^2 + (-2A + 2B + C)x + (4A + 2C) \end{aligned}$$

Comparing coefficients on both sides,

|                  |                         |                |
|------------------|-------------------------|----------------|
| $A + B = 0$      | $-2A + 2B + C = 0$      | $4A + 2C = 12$ |
| $1 + B = 0$      | $-2(1) + 2(-1) + C = 0$ | $C = 4$        |
| $\boxed{B = -1}$ |                         |                |

Putting the values of A, B and C in (i),

$$\frac{12}{(x+2)(x^2 - 2x + 4)} = \frac{1}{x+2} + \frac{-x+4}{x^2 - 2x + 4}$$

$$= \frac{1}{x+2} - \frac{x-4}{x^2-2x+4}$$

$$\begin{aligned} \int \frac{12}{(x+2)(x^2-2x+4)} dx &= \int \frac{d}{x+2} - \int \frac{x-1-3}{x^2-2x+4} dx \\ &= \ln|x+2| - \frac{1}{2} \int \frac{2x-2}{x^2-2x+4} dx + 3 \int \frac{dx}{x^2-2x+1+1^2+3} \\ &= \ln|x+2| - \frac{1}{2} \ln|x^2-2x+4| + 3 \int \frac{dx}{(x-1)^2+(\sqrt{3})^2} \\ &= \ln|x+2| - \frac{1}{2} \ln|x^2-2x+4| + 3 \left[ \frac{1}{\sqrt{3}} \tan^{-1} \left( \frac{x-1}{\sqrt{3}} \right) \right] + C \\ &= \ln|x+2| - \frac{1}{2} \ln|x^2-2x+4| + \sqrt{3} \tan^{-1} \left( \frac{x-1}{\sqrt{3}} \right) + C \end{aligned}$$

- (b) Find equations of two parallel lines, perpendicular to  $2x - y + 3 = 0$  such that the product of the x- and y-intercepts of each is 3. (5)

**Ans**  $2x - y + 3 = 0$

$$-y = -2x - 3$$

$$y = 2x + 3$$

$$\therefore \text{Slope of required lines } = m = -\frac{1}{2}$$

Equation of required lines are:

$$y = mx + c$$

$$= -\frac{1}{2}x + c = \frac{-x + 2c}{2}$$

$$2y = -x + 2c$$

$$\text{or } x + 2y - 2c = 0 \quad (i)$$

For x-intercept, put  $y = 0$  in (i),

$$\therefore x + 0 - 2c = 0$$

$$x = 2c$$

For y-intercept, put  $x = 0$  in (i),

$$2y - 2c = 0$$

$$2y = 2c \Rightarrow y = c$$

As product of x and y-intercept = 3

$$(2c)(c) = 3$$

$$2c^2 = 3$$

$$c^2 = \frac{3}{2}$$

$$c = \pm \sqrt{\frac{3}{2}}$$

Putting it in (1),

$$\begin{aligned}x + 2y - 2\left(\pm \sqrt{\frac{3}{2}}\right) &= 0 \\ \Rightarrow x + 2y \pm \sqrt{6} &= 0 \\ \therefore x + 2y + \sqrt{6} &= 0 \quad \text{and} \quad x + 2y - \sqrt{6} = 0\end{aligned}$$

---

**Q.7.(a)** Evaluate the definite integral  $\int_{\pi/6}^{\pi/2} \frac{\cos x}{\sin x(2 + \sin x)} dx$ . (5)

---

**Ans**

$$\text{Let } \sin x = t \Rightarrow \cos x dx = dt$$

$$\text{Limits: Let } x = \frac{\pi}{6} \Rightarrow t = \sin \frac{\pi}{6} = \frac{1}{2}$$

$$\text{Let } x = \frac{\pi}{2} \Rightarrow t = \sin \frac{\pi}{2} = 1$$

$$\int_{\pi/6}^{\pi/2} \frac{\cos x dx}{\sin x(2 + \sin x)} = \int_{1/2}^1 \frac{dt}{t(2 + t)} \quad (\text{i})$$

Making partial fractions,

$$\frac{1}{t(2 + t)} = \frac{A}{t} + \frac{B}{2 + t} \quad (\text{ii})$$

$$1 = A(2 + t) + B(t) \quad (\text{iii})$$

For A, let  $t = 0$

$$1 = A(2 + 0)$$

$$2A = 1$$

$$\boxed{A = \frac{1}{2}}$$

For B, let  $2 + t = 0$

$$t = -2$$

Putting the value of t in eq. (iii),

$$1 = 0 - 2B \Rightarrow B = -\frac{1}{2}$$

Putting the value of A and B in eq. (ii),

$$\frac{1}{t(2 + t)} = \frac{1}{2t} - \frac{1}{2(2 + t)}$$

Putting it in eq. (i),

$$\begin{aligned}
 \int_{\pi/6}^{\pi/2} \frac{\cos x dx}{\sin x(2 + \sin x)} &= \int_{1/2}^1 \frac{dt}{t(2 + t)} \\
 &= \frac{1}{2} \int_{1/2}^1 \frac{dt}{t} - \frac{1}{2} \int_{1/2}^1 \frac{dt}{2+t} \\
 &= \frac{1}{2} [\ln |t|]_{1/2}^1 - \frac{1}{2} [\ln |2+t|]_{1/2}^1 \\
 &= \frac{1}{2} \left[ \ln(1) - \ln\left(\frac{1}{2}\right) \right] - \frac{1}{2} \left[ \ln(2+1) - \ln\left(2+\frac{1}{2}\right) \right] \\
 &= \frac{1}{2} [0 - \ln(2)^{-1}] - \frac{1}{2} \left[ \ln(3) - \ln\left(\frac{5}{2}\right) \right] \\
 &= \frac{1}{2} [\ln(2)] - \frac{1}{2} [\ln(3) - \ln(5) + \ln(2)] \\
 &= \frac{1}{2} [\ln(2) - \ln(3) + \ln(5) - \ln(2)] \\
 &= \frac{1}{2} \ln\left(\frac{5}{3}\right)
 \end{aligned}$$

(b) Minimize  $z = 2x + y$  subject to the constraints (5)

$$x + y \geq 3, \quad 7x + 5y \leq 35, \quad x \geq 0, \quad y \geq 0$$

**Ans**

$$x + y \geq 3 \quad (\text{i})$$

$$x + y = 3 \quad (\text{iii})$$

Putting  $x = 0$  in (iii),

$$0 + y = 3 \Rightarrow y = 3$$

$\therefore (0, 3)$  is a point on (iii).

Putting  $y = 0$  in (iii),

$$x + 0 = 3 \Rightarrow x = 3$$

$\therefore (3, 0)$  is another point on (iii).

Putting  $x = 0, y = 0$  in (i),

$$0 + 0 > 3$$

$$0 > 3$$

Which is false. Hence solution region of (i) does not lie on the origin-side of (i).-

$$7x + 5y \leq 35 \quad (\text{ii})$$

$$7x + 5y = 35 \quad (\text{iv})$$

Putting  $y = 0$  in (iv),

$$0 + 5y = 35 \Rightarrow y = 7$$

$\therefore (0, 7)$  is a point on (iv).

Putting  $x = 0$  in (iv),

$$7x + 0 = 35 \Rightarrow x = 5$$

$\therefore (5, 0)$  is another point on (iv).

Putting  $x = 0, y = 0$  in (ii),

$$0 + 0 < 35$$

$$\therefore 0 < 35$$

Which is true. Hence solution region of (ii) lies on the origin-side of (ii).

Q.8.(a) Find equation of the line through the point  $(2, -9)$  and intersection of the lines (5)

$$2x + 5y - 8 = 0$$

$$3x - 4y - 6 = 0$$

**Ans**

$$2x + 5y - 8 = 0 \quad (i)$$

From (i),

$$2x = 8 - 5y$$

$$x = \frac{8 - 5y}{2} \quad (iii)$$

$$3x - 4y - 6 = 0 \quad (ii)$$

Putting eq. (iii) to eq. (ii),

$$3\left(\frac{8 - 5y}{2}\right) - 4y = 6$$

$$\frac{24 - 15y - 8y}{2} = 6$$

$$-23y + 24 = 12$$

$$-23y = -12$$

$$y = \frac{12}{23}$$

Putting the value of y in eq. (iii),

$$x = \frac{8 - 5\left(\frac{12}{23}\right)}{2} = \frac{\frac{184 - 60}{23}}{2}$$

$$= \frac{124}{46} = \frac{62}{23}$$

$\therefore \left(\frac{62}{23}, \frac{12}{23}\right)$  is the point of intersection of (i) and (ii).

Equation of line through  $(2, -9)$  and  $\left(\frac{62}{23}, \frac{12}{23}\right)$  is

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1} (x - x_1)$$

$$y + 9 = \frac{\frac{12}{23} + 9}{\frac{62}{23} - 2} (x - 2)$$

$$= \frac{\frac{12 + 207}{23}}{\frac{62 - 46}{23}} (x - 2)$$

$$y + 9 = \frac{219}{16} (x - 2)$$

$$16y + 144 = 219x - 438$$

$$219x - 16y - 582 = 0$$

(b) Show that the circles  $x^2 + y^2 + 2x - 2y - 7 = 0$  and  $x^2 + y^2 - 6x + 4y + 9 = 0$  touch externally. (5)

|  |   |
|--|---|
| <b>Ans</b> $x^2 + y^2 + 2x - 2y - 7 = 0$<br>$x^2 + y^2 + 2(1)x + 2(-1)y + (-7) = 0$<br>$g_1 = 1, f_1 = -1, c_1 = -7$<br>$c'_1(-g_1, -f_1) = c'_1(-1, 1)$<br>$r_1 = \sqrt{g_1^2 + f_1^2 - c_1}$<br>$= \sqrt{(1)^2 + (-1)^2 - (-7)}$<br>$= \sqrt{1 + 1 + 7} = \sqrt{9} = 3$<br>$ c'_1 c'_2  = r_1 + r_2 = 3 + 2 = 5$ | $x^2 + y^2 - 6x + 4y + 9 = 0$<br>$x^2 + y^2 + 2(-3)x + 2(2)y + 9 = 0$<br>$g_2 = -3, f_2 = 2, c_2 = 9$<br>$c'_2(-g_2, -f_2) = c'_2(3, -2)$<br>$r_2 = \sqrt{g_2^2 + f_2^2 - c_2}$<br>$= \sqrt{(-3)^2 + (2)^2 - 9}$<br>$= \sqrt{9 + 4 - 9} = \sqrt{4} = 2$ |
|--|---|

Also,  $|c'_1 c'_2| = \sqrt{(3 + 1)^2 + (-2 - 1)^2}$   
 $= \sqrt{16 + 9} = \sqrt{25} = 5$  (ii)

From (i) and (ii), it is prove that the given two circles touch externally.

**Q.9.(a)** Find an equation of the ellipse having foci  $(\pm 5, 0)$  and passing through the point  $\left(\frac{2}{3}, \sqrt{3}\right)$ . (5)

**Ans**  $F(\sqrt{5}, 0), F'(-\sqrt{5}, 0)$  through  $\left(\frac{3}{2}, \sqrt{3}\right)$

$$a_e = \sqrt{5} \Rightarrow e = \frac{\sqrt{5}}{a}$$

$$e^2 = \frac{a^2 - b^2}{a^2} \Rightarrow \frac{5}{a^2} \frac{a^2 - b^2}{a^2}$$

$$= a^2 - b^2 = 5 \Rightarrow a^2 = 5 + b^2 \quad (i)$$

Equation of required ellipse is:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \quad (ii)$$

Since  $\left(\frac{3}{2}, \sqrt{3}\right)$  lie on it, therefore

$$\frac{9}{4a^2} + \frac{3}{b^2} = 1 \Rightarrow \frac{9}{4a^2} + \frac{3}{b^2} = 1 \Rightarrow \frac{9b^2 + 12a^2}{4a^2b^2} = 1$$

$$\therefore 12a^2 + 9b^2 = 4a^2b^2 \quad (iii)$$

Putting the value of eq. (i) in eq. (iii),

$$12(5 + b^2) + 9b^2 = 4b^2(5 + b^2)$$

$$60 + 12b^2 + 9b^2 = 20b^2 + 4b^4$$

$$0 = 4b^4 - b^2 - 60$$

Let  $b^2 = t$

$$4t^2 - t - 60 = 0$$

$$t = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(4)(-60)}}{2 \times 4} = \frac{1 \pm \sqrt{1 + 960}}{8}$$

$$= \frac{1 \pm 31}{8} \Rightarrow \frac{1 + 31}{8} \text{ & } t = \frac{1 - 31}{8}$$

$$= \frac{32}{8} \quad = -\frac{30}{8}$$

$$t = 4, -\frac{15}{4} \quad \therefore -\frac{15}{4}$$

Gives imaginary roots, hence discard it.

$$\therefore b^2 = t = 4 \Rightarrow b = \pm 2$$

Putting the values in (i),

$$a^2 = 5 + 4 = 9 \Rightarrow a = \pm 3$$

Putting the values in (ii),

$$\frac{x^2}{9} + \frac{y^2}{4} = 1$$

- 
- (b) A particle acted upon by constant forces  $4\mathbf{i} + \mathbf{j} - 3\mathbf{k}$  and  $3\mathbf{i} - \mathbf{j} - \mathbf{k}$  is displaced from A(1, 2, 3) to B(5, 4, 1). Find the work done. (5)

**Ans** Let  $\vec{F}_1 = 4\mathbf{i} + \mathbf{j} - 3\mathbf{k}$  and  $\vec{F}_2 = 3\mathbf{i} - \mathbf{j} - \mathbf{k}$

$$\begin{aligned}\text{Total force} &= \vec{F} = \vec{F}_1 + \vec{F}_2 \\ &= (4\mathbf{i} + \mathbf{j} - 3\mathbf{k}) + (3\mathbf{i} - \mathbf{j} - \mathbf{k}) \\ &= 7\mathbf{i} + 0\mathbf{j} - 4\mathbf{k}\end{aligned}$$

$$\begin{aligned}\vec{d} &= \vec{AB} = (5 - 1)\mathbf{i} + (4 - 2)\mathbf{j} + (1 - 3)\mathbf{k} \\ &= 4\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}\end{aligned}$$

$$\begin{aligned}\text{Work done} &= \vec{F} \cdot \vec{d} = (7\mathbf{i} + 0\mathbf{j} - 4\mathbf{k})(4\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}) \\ &= (7 \times 4) + (0 \times 2) + (-4 \times -2) \\ &= 28 + 0 + 8 = 36\end{aligned}$$